

Development and validation of a modified QuEChERS protocol coupled to LC-MS/MS for simultaneous determination of multi-class antibiotic residues in honey

Hayam Lotfy, Amr H. Shendy, Medhat A. Al-Ghobashy, Sohair A. Gad Alla,

Abstract

LC-MS/MS assay was developed and validated according to EU guidelines for determination of nitrofuran metabolites and nitroimidazole residues in honey. Crude samples were acid-treated to liberate matrix-bound residues and a modified QuEChERS protocol was employed. Nitrofurantoin, furazolidone, furalfadone and nitrofurazone were determined via analysis of their metabolites AHD, AOZ, AMOZ and SEM, respectively while nitroimidazole residues; ronidazole (RNZ) and dimetridazole (DMZ) were determined directly. For all analytes, neat standard calibration curves, after correction for matrix effect were successfully employed. Decision limit (CC_a) and detection capability (CC_b) were below the MRPL for nitrofurans (1.00 µg kg⁻¹) and the recommended concentration for nitroimidazole (3.00 µg kg⁻¹), respectively. The CC_a, CC_b, percentage recovery and CV% ranges were 0.12-0.74 µg kg⁻¹, 0.21-1.27 µg kg⁻¹, 90.96-104.80% and 2.65-12.58%, respectively. This work is part of the national initiative for establishing a national monitoring program for drug residues in Egyptian honey.

Food Chemistry 2016, January